arXiv:hep-th/0408008v2 5 Aug 2004

Creation of a brane world with Gauss-Bonnet term

Koh-suke Aoyanagi* and Kei-ichi Maeda†
[1] *Department of Physics, Waseda University, 3-4-1 Okubo, Shinjuku-ku, Tokyo 169-8555, Japan*
[2] *Advanced Research Institute for Science and Engineering,*
Waseda University, Shinjuku, Tokyo 169-8555, Japan and
[3] *Waseda Institute for Astrophysics, Waseda University, Shinjuku, Tokyo 169-8555, Japan*

(Dated: February 1, 2008)

We study a creation of a brane world using an instanton solution. We analyze a brane model with a Gauss-Bonnet term in a bulk spacetime. The curvature of 3-brane is assumed to be closed, flat, or open. We construct instanton solutions with branes for those models, and calculate the value of the actions to discuss an initial state of a brane universe.

PACS numbers: 98.80.cq

I. INTRODUCTION

Although a big bang universe is very successful, it predicts the existence of an initial singularity. To resolve such a difficulty, we have to search for new gravitation theory such as quantum gravity. However, we could so far not find it. As a first step to reach it, we may consider a mini-super space and quantize the isotropic and homogeneous universe, which is the so-called quantum cosmology [1, 2]. Vilenkin claimed that the universe is created from nothing [1]. This approach is based on the picture that the universe is spontaneously nucleated in a de Sitter space. The mathematical description of this nucleation is analogous to a quantum tunneling through a potential barrier [3]. Another approach to quantum cosmology is developed by Hartle and Hawking [2], who proposed that the wave function of the Universe is given by a path integral over non-singular compact Euclidean geometries, which is called a “no boundary” boundary condition.

When we discuss the early stage of the universe, however, a unified theory of fundamental interactions and particles will play a very important role. Among such unified theories, a string/M theory is the most promising candidate, which is constructed in higher-dimensional spacetime. Based on such higher-dimensional theories, new cosmological scenario has been proposed, that is a brane cosmology. The prototype of a brane world was first discussed in [4, 5]. Combined it with the idea of D-brane found by Polchinski in a string theory [6], a new paradigm of a brane world has been developed [7]. Randall and Sundrum also proposed new brane models [8, 9]. In their second model [9], it was shown that the four-dimensional gravity is recovered even in an infinite bulk spacetime. It also provides new terms in the Friedmann equation of a brane universe [10]. The effective 4-dimensional (4D) Einstein equations with these new terms are obtained covariantly by projection of the 5-dimensional (5D) spacetime onto the brane world [11], by which we understand naturally the origin of new terms.

If we believe such higher-dimensional cosmological scenario, we have to invoke how such a universe is created. In particular, because a brane structure is highly inhomogeneous in a higher-dimensional bulk spacetime, we may wonder how such a brane universe is born and starts to evolve. As for creation of a brane universe, a few works has been so far done. Garriga and Sasaki first constructed an inflating brane instanton of Randall-Sundrum model [12]. This instanton is obtained by gluing two spherical parts of AdS$_5$. Hawking, Hetrog and Reall consider the creation using an instanton, and discuss inflation and a fluctuations during the de Sitter phase in the model which contains the quantum correction term called a trace anomaly on the brane [13, 14]. The trace anomaly term is first discussed by Starobinsky for inflation [15]. Another quantum correction term called induced gravity is introduced on the brane by Dvali, Gabadadze, and Porrati [16]. For this model, the effective gravitational equations on the brane are derived [17]. This may provide new de Sitter phase without a cosmological constant.

When we discuss such quantum effects on the brane, we may also have to include quantum effects in the bulk. In the higher dimensional theory, the higher curvature correction terms should be added to the Einstein-Hilbert action. These terms appear in the low energy effective action of string theory via quantum one-loop corrections. In fact the low energy effective actions of some sting theories include $\mathcal{R}^{ABCD}\mathcal{R}_{ABCD}$ interactions, but this term gives rise to a ghost. In order to resolve this problem, Zwiebach [18] (also see [19]) introduce a ghost-free Gauss-Bonnet combination. Hence, the Gauss-bonnet term should be included when we discuss a brane universe with some quantum corrections.

For RS II type model in the Einstein-Gauss-Bonnet theory, it is shown that the graviton zero mode is localized at low energies as in the original RS II model and that the correction of the Newton’s law becomes milder by including the Gauss-Bonnet term [20, 21]. Also the covariant effective 4D Einstein equation is given by Maeda and Torii [22].

In this paper we consider the creation of a brane uni-

---

*E-mail address : aoyanagi@gravity.phys.waseda.ac.jp
†E-mail address : maeda@gravity.phys.waseda.ac.jp

verse using an instanton solution in the Einstein-Gauss-Bonnet theory. We construct a brane instanton including a Gauss-Bonnet term. In §II, we present Euclidean action and Euclidean equations of motion in the Einstein-Gauss-Bonnet theory. In §III, we obtain the instanton solutions which 3-curvature is positive, zero or negative. We show that only for a closed model with positive curvature there exist instanton solutions either with a single brane or with two branes. For a flat or an open model, there does not exist any instanton solution with a single brane because of a boundary condition. We also calculate the value of the Euclidean actions to discuss an initial state of a brane universe. In Section IV we consider the model including a trace anomaly term on the brane. We show there exists instanton solution with an inflating brane. In §V, conclusion follows.

II. ACTION AND EQUATIONS OF MOTION

The Euclidean action for the brane world with Gauss-Bonnet term in the 5D space time is described by two parts: One in a bulk spacetime ($\mathcal{M}$) and the other in brane boundary hypersurfaces ($\partial\mathcal{M} = \sum_i \partial\mathcal{M}_i$), i.e.

$$S_E = S_E^{\text{bulk}} + S_E^{\text{brane}}\,. \tag{2.1}$$

The bulk action is given by

$$S_E^{\text{bulk}} = -\frac{1}{2\kappa_5^2}\int_{\mathcal{M}} dx^5\sqrt{g}\left[\mathcal{R} - 2\Lambda + \alpha\mathcal{L}_{GB}\right], \tag{2.2}$$

where

$$\mathcal{L}_{GB} = \mathcal{R}^2 - 4\mathcal{R}_{AB}\mathcal{R}^{AB} + \mathcal{R}_{ABCD}\mathcal{R}^{ABCD}. \tag{2.3}$$

κ_5^2 is the 5D gravitational constant, Λ is a cosmological constant, $\mathcal{R}, \mathcal{R}_{AB}$ and $\mathcal{R}_{ABCD}$ are the 5D scalar curvature, Ricci tensor and Riemann tensor, respectively, and α is a constant, which is related to a string coupling constant. The induced 4D metric $h_{\mu\nu}$ on a 3-brane is defined by

$$h_{AB} = g_{AB} - n_A n_B, \tag{2.4}$$

where n_A is the spacelike unit vector field which normal to the brane hypersurface. The action of the branes is given by the following form:

$$S_E^{\text{brane}} = -\sum_i \int_{\partial\mathcal{M}_i} d^4x\sqrt{h}\left[\frac{1}{\kappa_5^2}L_{\text{surface}}(\partial\mathcal{M}_i) - \lambda_i\right], \tag{2.5}$$

where

$$L_{\text{surface}} = K + 2\alpha(J - 2G^{\rho\sigma}K_{\rho\sigma}) \tag{2.6}$$

is a surface term of the 5D gravitational action [25, 26, 27]. λ_i is a tension on the i-th brane, $K_{\mu\nu}$ is the extrinsic curvature of a brane, $K = K^\mu_\mu$, and $G_{\mu\nu}$ is the Einstein tensor of the induced metric $h_{\mu\nu}$. J is a trace of $J_{\mu\nu}$, which is given by some combination of the extrinsic curvature defined by

$$\begin{aligned} J_{\mu\nu} &= \frac{1}{3}\left(2KK_{\mu\rho}K^\rho{}_\nu + K_{\rho\sigma}K^{\rho\sigma}K_{\mu\nu}\right. \\ &\qquad \left.-2K_{\mu\rho}K^{\rho\sigma}K_{\sigma\nu} - K^2K_{\mu\nu}\right). \end{aligned} \tag{2.7}$$

The total action ($S_E = S_E^{\text{bulk}} + S_E^{\text{brane}}$) gives the field equations as

$$\mathcal{G}_{AB} + \alpha\mathcal{H}_{AB} = -\Lambda g_{AB} - \sum_i \lambda_i g_{AB}\delta(\partial\mathcal{M}_i), \tag{2.8}$$

where

$$\mathcal{G}_{AB} = \mathcal{R}_{AB} - \frac{1}{2}g_{AB}\mathcal{R}, \tag{2.9}$$

and

$$\begin{aligned} \mathcal{H}_{AB} &= 2\left[\mathcal{R}\mathcal{R}_{AB} - 2\mathcal{R}_{AC}\mathcal{R}^C_{\ B} - 2\mathcal{R}^{CD}\mathcal{R}_{ACBD}\right. \\ &\qquad \left.+\mathcal{R}_A^{\ CDE}\mathcal{R}_{BCDE}\right] - \frac{1}{2}g_{AB}\mathcal{L}_{GB}. \end{aligned} \tag{2.10}$$

Since we are looking for an instanton solution, we assume a highly symmetric Euclidean spacetime, which metric is given by

$$ds_E^2 = dr^2 + b(r)^2\gamma_{\mu\nu}dx^\mu dx^\nu, \tag{2.11}$$

where $\gamma_{\mu\nu}$ is a 4D metric with maximal symmetry. This maximally symmetric 4D space is classified into three cases by the signature of curvature, i.e. k =0 (zero), 1 (positive), or -1 (negative). It corresponds to the curvature sign of the Friedmann universe after creation. Since the Euclidian space must be compact when we discuss its creation, in the case of k=0 or -1, we have to make a space compact by identification. Then the flat spacetime is a 4D torus, and that with $k = -1$ has more complicated topology. Although the spacetimes are compact, we shall call them "flat" and "open" for $k = 0$ and -1 as well as "closed" for $k = 1$.

The equations of motion under the above ansatz are given by

$$\begin{aligned} 3\left(\frac{b''}{b} - \frac{k - b'^2}{b^2}\right) + 12\alpha\frac{(k - b'^2)}{b^3}b'' \\ = -\Lambda - \sum_i \bar{\lambda}_i\delta(r - r_i), \end{aligned} \tag{2.12}$$

$$6\frac{(k - b'^2)}{b^2}\left\{1 + 2\alpha\frac{(k - b'^2)}{b^2}\right\} = \Lambda, \tag{2.13}$$

where the prime denotes the derivative with respect to r and $\bar{\lambda}_i = \kappa_5^2\lambda_i$.

By integrating the first equation for a small interval $(r_i - \epsilon, r_i + \epsilon)$ including a brane, one obtain the junction condition [28] at $r = r_i$ as

$$\left[\frac{b'}{b}\left\{3 - 4\alpha\left(\frac{b'^2}{b^2} - \frac{3k}{b^2}\right)\right\}\right]^{\pm} = -\bar{\lambda}_i\,, \tag{2.14}$$



(a)

(b)

FIG. 1: brane instanton. The thick vertical circle at $r = r_i$ represents the 4-sphere brane at which the two identical 5-dimensional anti-de Sitter spaces are glued.

where $[\ ]^{\pm}$ denotes

$$[A]^{\pm} = A(r_i+\epsilon) - A(r_i-\epsilon) \equiv A_+ - A_- \qquad (2.15)$$

With the ansatz of Z_2 symmetry, which gives the relation of $[A]^{\pm} = 2A_+ = -2A_-$, we obtain

$$\frac{b'}{b}\left\{3 - 4\alpha\left(\frac{b'^2}{b^2} - \frac{3k}{b^2}\right)\right\} = \mp\frac{\bar{\lambda}_i}{2}, \qquad (2.16)$$

Here the upper (lower) sign is applied at $r = r_2$ (at $r = r_1$) for a two-brane model. For a single brane model, we apply the upper case at $r = r_0$. Through this paper, we use the notation $r_i (i = 0, 1, 2)$ where i means the number of a brane. For a two-brane model $i = 1$ and 2 stand for a negative and a positive tension brane, respectively. While for a single brane model, we use $i = 0$ standing for a brane (see Fig. 1).

With equations of motion the action is reduced to

$$S_E = -\frac{8V_4^{(k)}}{\kappa_5^2}\int dr\left\{\Lambda + 2\frac{b''}{b} - 3\frac{k-b'^2}{b^2}\right\} + \frac{V_4^{(k)}}{\kappa_5^2}\sum_i\left[\frac{b'}{b}\left\{1 - 12\alpha\left(\frac{b'^2}{b^2} - \frac{3k}{b^2}\right)\right\}\right]^{\pm}_{r=r_i} \qquad (2.17)$$

where $V_4^{(k)}$ is a volume of 4D manifold with the curvature k.

III. INSTANTON SOLUTION

We first provide a solution of Eqs. (2.12) and (2.13) in a bulk. Eq. (2.13) gives the quadratic equation,

$$6X + 12\alpha X^2 = \Lambda, \qquad (3.1)$$

where

$$X \equiv \frac{k - b'^2}{b^2}. \qquad (3.2)$$

If $\alpha \neq 0$, Eq. (3.1) gives two solutions;

$$X = X_{\pm} \equiv \frac{-1 \pm \eta}{4\alpha}, \qquad (3.3)$$

where $\eta = \sqrt{1 + 4\alpha\Lambda/3}$. Here we find two solutions; one is called as a plus-branch and the other is a minus-branch. The limit of $\alpha \to o$ exists only for the plus-branch solution.

In this paper we consider only a negative cosmological constant ($\Lambda < 0$). [1] Then we have the following constraint,

$$-\frac{3}{4\alpha} \leq \Lambda < 0\,, \qquad (3.4)$$

which is required in order that $X_{\pm}$ is a real value. The range of η is restricted as $0 \leq \eta < 1$ from the constraint (3.4). Under this condition, $X_{\pm}$ is always negative. We then introduce a typical length scale as

$$l_{\pm} \equiv (-X_{\pm})^{-1/2} = \left[-\frac{3(1 \pm \eta)}{\Lambda}\right]^{1/2}. \qquad (3.5)$$

Since $X' = 0$, we find $b'' = l_{\pm}^2 b$ from Eq. (3.2). This with Eq. (3.1) guarantee Eq. (2.12). Hence Eq. (3.3) gives a bulk solution.

In what follows, we discuss the instanton solutions for each value of k in order.

A. de Sitter brane instanton ($k = 1$)

In the case of $k = 1$, the solution (3.3) is written by,

$$b(r) = l_{\pm}\sinh\left(\frac{r}{l_{\pm}}\right), \qquad (3.6)$$

which also satisfies Eq. (2.12) in a bulk. From this bulk solution, we construct an instanton solution by cutting the space at $r = r_i$ and gluing two copies of it on the surfaces of the corresponding point in order that a compact Euclidean manifold (instanton) is obtained. At $r = r_i$, we impose the Israel's junction condition (2.16) with $k = 1$. For a single brane instanton, we impose "no boundary boundary condition" at the origin [2]. For a two-brane model, we impose the junction condition at r_1 and r_2. As a result, the tension of i-th brane is determined by these junction conditions. Substituting (3.3) into (2.16),

---

[1] In the case of $\Lambda = 0$, there also exist instanton solutions. For $k = 1$, the minus-branch solution is included in (3.3), and the plus-branch solution is also included in the limit of $X_+ \to 0$. This plus-branch solution is 5D Milne universe. For $k = 0$ the solutions of both branches is included in (3.15). For $k = -1$ only the minus-branch solution exists and is included in (3.21).

$$\bar{\lambda}_i^{(\pm)} = (-1)^i \frac{2}{l_\pm}\left[(2\pm\eta)\frac{\cosh(r_i/l_\pm)}{\sinh(r_i/l_\pm)} + 2(1\mp\eta)\left(\frac{\cosh(r_i/l_\pm)}{\sinh^3(r_i/l_\pm)}\right)\right] . \quad (3.7)$$

When we take the limit of $\alpha \to 0$ in the plus-branch, we recover the Garriga-Sasaki instanton, i.e. $l_\pm = l \equiv \sqrt{-6/\Lambda}$, $\eta = 1$, and

$$\bar{\lambda}_i = (-1)^i \frac{6}{l}\coth\left(\frac{r_i}{l}\right) . \quad (3.8)$$

Here we note a 'critical' tension. In a brane model with the Gauss-Bonnet term in a bulk, we find some contribution from the Gauss-Bonnet term in a 4-dimensional cosmological constant. As a result, the fine-tuned value of the tension to find the 4D Minkowski brane, which we shall call a critical tension, is modified from the Randall-Sundrum's value. The fine-tuned value is given by [22]

$$\alpha\bar{\lambda}_{\rm cr}^2 = 1 - 4\alpha\Lambda \mp \left(1 + \frac{4}{3}\alpha\Lambda\right)^3 . \quad (3.9)$$

In our case, if we take the limit of $r \to \infty$, the brane approaches to the 4D Euclidian flat space because the radius of the brane ($b(r)$) becomes infinitely large and the curvature of the brane (S^4 manifold) vanishes. In this limit, the tension of the brane (3.8) is

$$\bar{\lambda}_{i,\rm cr} = (-1)^i \frac{2}{l_\pm}(2\pm\eta). \quad (3.10)$$

This value is consistent with the above generalized Randall-Sundrum tuning condition (3.9). Using this critical tension, the tension of a positive-tension brane is divided into two parts $\bar{\lambda} = \bar{\lambda}_{\rm cr} + \triangle\bar{\lambda}$. It turns out that $\triangle\bar{\lambda}$ is always positive because $\bar{\lambda}$ decreases monotonically with respect to r for $0 \le \eta < 1$ (see Fig. 2). Hence, this brane has always a positive effective cosmological constant, that is, de Sitter brane.

We calculate the action in order to discuss which state is most plausible when the brane universe is created. The total Euclidian action for this solution is calculated as

$$S_E = -\frac{V_4^{(+)}}{\kappa_5^2} l_\pm^3 \left\{\left[l_\pm^3(\Lambda l_\pm^2 + 9)\sinh\left(\frac{r_2}{l_\pm}\right)\cosh\left(\frac{r_2}{l_\pm}\right) + 3l_\pm^3(\Lambda l_\pm^2 + 5)\left(\frac{r_2}{l_\pm}\right)\right] - [r_2 \to r_1]\right\}, \quad (3.11)$$



(a)

(b)

FIG. 2: The tension for a de Sitter brane instanton with $\eta = 0, 1/3, 2/3$, and 1. The figures (a) and (b) correspond to the cases of a plus and a minus branches, respectively. $d\bar{\lambda}/dr$ is always negative because $2\pm\eta > 0$ and $1\mp\eta > 0$ for the range of $0 \le \eta < 1$. Since $\bar{\lambda}$ monotonically decreases with respect to r and approaches the critical tension as r gets large, $\triangle\bar{\lambda}$ is always positive.

where we have used the relation $\Lambda l_\pm^4 + 6l_\pm^2 - 12\alpha = 0$. Here $V_4^{(+)} = 8\pi^2/3$ because of S^4 manifold. We rewrite the action using the variable η,

$$S_E = -\frac{3V_4^{(+)}}{\kappa_5^2} l_\pm^3 \left\{\left[(2\mp\eta)\sinh\left(\frac{r_2}{l_\pm}\right)\cosh\left(\frac{r_2}{l_\pm}\right) + (2\mp 3\eta)\left(\frac{r_2}{l_\pm}\right)\right] - [r_2 \to r_1]\right\}. \quad (3.12)$$

For single brane instanton, the action is given by replacing r_2 and r_1 with r_0 and 0, respectively.

Those actions do not have any minimum value, and get small when the distance between two branes or the size of the brane becomes large. Although we can claim that the brane universe may be created as large as possible, we cannot predict the initial size.

The evolution of a brane after creation is given by analytic continuation of the Euclidean metric

$$ds_E^2 = dr^2 + l^2 \sinh^2(r/l_\pm)\left(d\chi^2 + \sin^2\chi d\Omega_{(3)}^2\right) \quad (3.13)$$

by the Wick rotation, where $d\Omega_{(3)}^2$ is the metric of the 3-sphere. It is done by substituting $\chi \to iHt + \pi/2$, which leads to

$$\begin{aligned} ds^2 &= dr^2 + (l_\pm H)^2 \sinh(r/l_\pm) \\ &\quad \times[-dt^2 + H^{-2}\cosh^2(Ht)d\Omega_{(3)}^2], \end{aligned} \quad (3.14)$$

where $H \equiv l_\pm \sinh(r_i/l_\pm)$ ($r_i = r_0$ or r_2) is the radius of a brane. After the creation of this spacetime, the universe inflates. If $\triangle\bar{\lambda}$ is given by some potential of a scalar field and will decrease to zero, inflation will end (see §. IV).

B. flat brane instanton ($k = 0$)

In the case of a flat brane ($k = 0$), the solution of Eq. (3.2) is $b(r) = b_0 e^{\pm r/l_\pm}$, where b_0 is an integrating constant. Due to the Z_2 symmetry, we consider only the plus sign without loss of generality, i.e.

$$b(r) = b_0 e^{r/l_\pm}\,. \quad (3.15)$$

We can construct an instanton solution in the same way as the previous case. We can impose the junction conditions at the brane boundaries ($r = r_1$ and r_2, or r_0). This solution, however, does not satisfy no-boundary boundary condition, because $b(r)$ dose not vanish at any point r. Thus we cannot construct a single brane instanton solution. Here we consider only a two-brane model.

The tension of i-th brane is determined by the junction condition (2.16) as

$$\frac{b'}{b}\left(3 - 4\alpha\frac{b'^2}{b^2}\right) = \mp\frac{\bar{\lambda}_i}{2}. \quad (3.16)$$

Substituting Eq. (3.15) into Eq. (3.16),

$$\bar{\lambda}_i^{(\pm)} = (-1)^i \frac{2}{l_\pm}(2 \pm \eta)\,. \quad (3.17)$$

This tension is independent of the position of a brane and is the same as the critical tension (3.10).

As for the Euclidean action of this solution, we find

$$S_E = -\frac{2V_4^{(0)} b_0^4 (2 \mp 3\eta)}{\kappa_5^2\, l_\pm}\left[e^{4r_2/l_\pm} - e^{4r_1/l_\pm}\right]. \quad (3.18)$$

Here $V_4^{(0)}$ is the volume of 4D torus. This action does also not have any minimum value with respect to the distance between two branes. Then we cannot predict the initial size of the universe. Note that we can obtain one-brane RS II model in the limit of $r_1 \to -\infty$. In that case, the total action S_E is still finite. Therefore, an instanton solution with one flat brane exists.

The evolution of a brane universe after creation is also given by analytic continuation of the Euclidean metric

$$ds_E^2 = dr^2 + b_0^2 e^{2r/l_\pm}\left(d\tau^2 + dx^2 + dy^2 + dz^2\right) \quad (3.19)$$

by the Wick rotation. Substituting $\tau \to it$, we obtain

$$ds^2 = dr^2 + b_0^2 e^{2r/l_\pm}[-dt^2 + dx^2 + dy^2 + dz^2]\,. \quad (3.20)$$

We recover the 4D Minkowski spacetime.

C. anti de Sitter brane instanton ($k = -1$)

In the case of an 'open' brane model ($k = -1$), the solution of Eq. (3.2) is given as

$$b(r) = l_\pm \cosh\left(\frac{r}{l_\pm}\right)\,. \quad (3.21)$$

We impose the junction condition at the boundary ($r = r_1$ and r_2). As the same reason for a flat brane model, this solution does not provide a single brane model.

The tension of i-th brane is determined by the junction condition (2.16) as

$$\frac{b'}{b}\left(3 - 4\alpha\frac{b'^2}{b^2} - 12\alpha\frac{1}{b^2}\right) = \mp\frac{\bar{\lambda}_i}{2}. \quad (3.22)$$

Substituting Eq. (3.21) into Eq. (3.22), we obtain

$$\begin{aligned} \bar{\lambda}_i^{(\pm)} &= (-1)^i \frac{2}{l_\pm}\left[(2 \pm \eta)\frac{\sinh(r_i/l_\pm)}{\cosh(r_i/l_\pm)}\right. \\ &\qquad \left. - 2(1 \mp \eta)\frac{\sinh(r_i/l_\pm)}{\cosh^3(r_i/l_\pm)}\right]. \end{aligned} \quad (3.23)$$

In the limit of $r \to \infty$, we recover the critical tension (3.10), since the curvature of a brane vanishes. Furthermore, if we divide the tension (3.23) into two parts; $\bar{\lambda} = \bar{\lambda}_{\rm cr} + \triangle\bar{\lambda}$, we find that $\triangle\bar{\lambda}$ is always negative for $0 \le \eta < 1$. Since this tension gives a negative effective cosmological constant on the brane, the brane is anti-de Sitter spacetime.

The total action ($S_E = S_E^{\rm bulk} + S_E^{\rm brane}$) is given by

$$\begin{aligned} S_E &= -\frac{3V_4^{(-)}}{\kappa_5^2} l_\pm^3 \left\{\left[(-2 \pm \eta)\cosh\left(\frac{r_2}{l_\pm}\right)\sinh\left(\frac{r_2}{l_\pm}\right)\right.\right. \\ &\qquad \left.\left. + (2 \mp 3\eta)\left(\frac{r_2}{l_\pm}\right)\right] - [r_2 \to r_1]\right\} \end{aligned} \quad (3.24)$$

Here $V_4^{(-)}$ is the volume of 4D manifold with $k = -1$. Again we do not have any minimum in this action. We cannot give any prediction for a created brane spacetime.

To discuss the evolution of the brane universe after creation, we have to perform analytically continuation of the Euclidian space;

$$ds_E^2 = dr^2 + l_\pm^2 \cosh^2(r/l_\pm) ds_{E,4}^2 \quad (3.25)$$



(a)

(b)

FIG. 3: The tension for an open brane instanton with $\eta = 0, 1/3, 2/3$, and 1. Figs. (a) and (b) depict the tension of a plus and a minus branches, respectively. $d\bar{\lambda}/dr$ changes sign from negative to positive and $\bar{\lambda}$ approaches the critical tension in the limit of $r \to \infty$. Since $\bar{\lambda} = 0$ at $r = 0$, $\triangle\bar{\lambda}$ is always negative.

with

$$ds^2_{E,4} = d\chi^2 + \sinh^2\chi\left(d\psi^2 + \sin^2\psi d\Omega^2_{(2)}\right) . \quad (3.26)$$

By the double Wick rotations, i.e. $\chi \to i(t + \pi/2)$ and $\psi \to i\phi$ [29], we obtain the metric of the brane as

$$ds^2_4 = -dt^2 + \cos^2 t\left(d\phi^2 + \sinh^2\phi d\Omega^2_{(2)}\right) , \quad (3.27)$$

which gives the AdS spacetime.

IV. TRACE ANOMALY

In the previous section, although we can find instanton solutions for several situations, we cannot provide the most plausible state of the brane universe because the action does not give any minimum value. If we choose the critical tension to obtain zero cosmological constant, only a flat brane instanton is possible. The size of the brane universe, however, is not determined. Any distance between two brane is possible. We again loose the predictability. In this section we discuss another effect which may give a prediction of the initial state of the brane universe. In a curved space time, we know that even in the absence of classical gravitational action, quantum fluctuation of matter fields provides a nontrivial gravitational action through a trace anomaly term $\langle\tau_{\mu\nu}\rangle$. In the case of free, massless, and conformally invariant fields, these quantum corrections take a simple form [30]. Those term were discussed in the context of inflationary scenario [15] and creation of the universe [13, 31]. In [13], assuming the critical tension, the size of the brane universe is fixed. Then, in the present case, we also take into account of the contribution of the trace anomaly, our junction condition is modified as

$$\begin{aligned}[B_{\mu\nu}]^{\pm} &\equiv \left[-\frac{b'}{b}\left\{3 - 4\alpha\left(\frac{b'^2}{b^2} - 3\frac{k}{b^2}\right)\right\}\right]^{\pm} h_{\mu\nu} \\ &= -\kappa_5^2\left(\tau_{\mu\nu} + \langle\tau_{\mu\nu}\rangle\right) , \end{aligned} \quad (4.1)$$

where $\tau_{\mu\nu}$ is the energy-momentum tensor of the brane matter fields and $\langle\tau_{\mu\nu}\rangle$ is its trace anomaly term including the tension of the brane, which is given by

$$\langle\tau_{\mu\nu}\rangle = -\lambda h_{\mu\nu} + H^{(1)}_{\mu\nu} + H^{(3)}_{\mu\nu} . \quad (4.2)$$

$H^{(1)}_{\mu\nu}$ and $H^{(3)}_{\mu\nu}$ take the following forms;

$$\begin{aligned} H^{(1)}_{\mu\nu} &= -k_1\left(2RR_{\mu\nu} - \frac{1}{2}h_{\mu\nu}R^2 - 2\nabla_\mu\nabla_\nu R \right. \\ &\quad \left. +2h_{\mu\nu}\nabla^\alpha\nabla_\alpha R\right) , \end{aligned} \quad (4.3)$$

$$\begin{aligned} H^{(3)}_{\mu\nu} &= k_3\left(-R_\mu{}^\sigma R_{\nu\sigma} + \frac{2}{3}RR_{\mu\nu} + \frac{1}{2}h_{\mu\nu}R^{\sigma\tau}R_{\sigma\tau} \right. \\ &\quad \left. -\frac{1}{4}h_{\mu\nu}R^2\right) , \end{aligned} \quad (4.4)$$

where R and $R_{\mu\nu}$ are the 4D scalar curvature and Ricci tensor respectively. The coefficient k_1 may not appear in $\mathcal{N} = 4$ super conformal Yang-Mills theory but can be included to obtain a successful inflationary scenario. While k_3 is uniquely determined:

$$k_3 = \frac{1}{2880\pi^2}(2N_0 + 11N_{1/2} + 62N_1), \quad (4.5)$$

where N_0, $N_{1/2}$, and N_1 are the number of quantum fields with spins 0, 1/2, and 1, respectively.

We shall include the trace anomaly terms for our instanton solutions. By using the metric (2.11),

$$R = \pm\frac{12}{b^2}, \quad R_{\mu\nu} = \pm\frac{3}{b^2}h_{\mu\nu}, \quad \nabla_\mu R = 0, \quad (4.6)$$

where the upper (or lower) sign corresponds to the $k = 1$ brane instanton (or $k = -1$ brane one). For a flat brane, the scalar curvature and Ricci tensor vanish.

The trace anomaly terms are then given by

$$H^{(1)}_{\mu\nu} = 0, \tag{4.7}$$

$$H^{(3)}_{\mu\nu} = -\frac{3k_3}{b^4} h_{\mu\nu}. \tag{4.8}$$

We obtain

$$\langle \tau_{\mu\nu} \rangle = -\frac{3k_3}{b^4} h_{\mu\nu}. \tag{4.9}$$

Hence the junction condition is

$$\left[\frac{b'}{b}\left\{3 - 4\alpha\left(\frac{b'^2}{b^2} - \frac{3k}{b^2}\right)\right\}\right]^{\pm} = -\left(\bar{\lambda} + \frac{3\bar{k}_3}{b^4}\right), \tag{4.10}$$

where $\bar{\lambda} = \kappa_5^2 \lambda$ and $\bar{k}_3 = \kappa_5^2 k_3$.

For a positive tension brane (either a single brane or the second brane of two-brane model), we obtain

$$\bar{\lambda} = 2\frac{b'}{b}\left\{3 - 4\alpha\left(\frac{b'^2}{b^2} - \frac{3k}{b^2}\right)\right\} - \frac{3\bar{k}_3}{b^4}. \tag{4.11}$$

Since a trace anomaly (4.9) is always negative (or zero for a flat brane), the tension is always below the value obtained previously. It turns out that only de Sitter brane with $k = 1$ is possible if we adopt the critical tension. We show the tension in terms of including trace anomaly term in Fig. 4.

$$\bar{\lambda}^{(\pm)} = \frac{2}{l_\pm}\left\{(2 \pm \eta) + \frac{2(1 \mp \eta)}{\sinh^2(r_i/l_\pm)}\right\} - \frac{3\bar{k}_3}{l_\pm^4 \sinh^4(r_i/l_\pm)}. \tag{4.12}$$

If the tension is the critical value (3.10), we find a unique solution with a finite radius.

For a two-brane model, if the tension on one brane is critical, the other is not the case. For example, if the tension of the positive tension brane (r_2) is critical, a effective cosmological constant on the negative tension brane (r_1) is positive. Conversely, if the tension of the negative tension brane (r_1) is critical, we find AdS universe on the positive-tension brane. For a single brane model, the radius of the created brane universe is fixed. In Fig. 5, we depict the ratio of the size of a single brane with respect to η to that without the Gauss-Bonnet term, which is obtained by [13]. We find that the difference is not so large. For the plus branch, the radius of the created universe is always smaller than that without Gauss-Bonnet term. While, for negative branch, it highly depends on the value of η. If we take the limit of $\eta \to 1 (\alpha \to 0)$, the radius gets much larger.

Note that since the trace anomaly always vanishes for a flat brane, the flat two-brane model at any distance is possible.



(a)

(b)

FIG. 4: The tension for a de Sitter brane instanton with $\eta = 2/3$ and $\bar{k}_3/l_\pm = 3, 10, 10^2, 10^3$, and 0, where $\bar{k}_3 = 0$ gives the case without the trace anomaly term. The figures (a) and (b) correspond to the cases of a plus- and a minus-branches, respectively. Since $\lim_{r_i \to \infty}[\bar{\lambda} - \bar{\lambda}_{\rm cr}] = 0+$, we find a unique solution at finite radius if the tension is the critical value $\bar{\lambda}_{\rm cr}$.

V. CONCLUSION

We have presented instanton solutions in the model with a Gauss-Bonnet term. If a brane is a closed universe, we find both two-brane and single-brane de Sitter brane instantons. For a flat brane universe, we can also construct a two-brane instanton solution. As for a single-brane model, a compact bulk spacetime with a single brane is not possible because a flatness of the brane is not consistent with a no-boundary boundary condition. However, the RS II type instanton with a non-compact bulk spacetime is allowed because its Euclidean action is finite. For an anti-de Sitter brane instanton with negative curvature, we can also construct only a two-brane instanton solution. In this case, the RS II type instanton does not exist because the Euclidean action diverges.

Although we find instanton solutions in the model with



FIG. 5: The radius of a de Sitter brane instanton with respect to η. The figure shows the ratio of the nucleation radius to that without the Gauss-Bonnet term. The figures (a) and (b) correspond to the cases of a plus- and a minus- branches, respectively. Here R is the radius of the positive brane : $R \equiv l_\pm \sinh(r_i/l_\pm)$ and R_1 is the radius of the positive brane with $\eta = 1$: $R_1 \equiv l_1^\pm \sinh(r_i/l_1^\pm)$, where $l_1^\pm \equiv l_\pm(\eta = 1)$.

Gauss-Bonnet term, we cannot predict the initial state of the brane universe. This is because the Euclidian action has no minimum value. For a flat and an anti-de Sitter brane, the volume of a created brane universe $V_4^{(k)}$ is not fixed as well. $V_4^{(k)}$ can be arbitrary.

As for the tension of a brane, although the critical tension requires a fine-tuning. we need such a choice to explain the present small value of the universe. Such a tuning could be obtained in some super symmetric theories such as Horava-Witten model. Here we assume such a tuned value, i.e. the critical tension. Then we also include trace anomaly terms on the brane. In this case, we can predict the size of the universe. If we have a single-brane universe with no-boundary boundary condition, the created universe is in de Sitter phase and naturally evolves into inflationary stage.

According to the recent analysis by Charmousis and Dufaux [23], the models with a negative-tension brane show its instability in the Einstein-Gauss-Bonnet theory. However such a problem is absent in the non-compact RS II model. For de Sitter brane instanton with two branes, this instability may be smeared out by the existence of trace anomaly terms similarly to the inclusion of induced gravity term on the branes. Furthermore in order to predict the initial state of created brane universe, we need to include other important effects such as the Casimir energy, which are not taken into account here. These issues are left for a future study.

Acknowledgments

We would like to thank S. Mizuno, N. Okuyama and T. Torii for useful discussions and comments. This work was partially supported by the Grant-in-Aid for Scientific Research Fund of the MEXT (No. 14540281) and by the Waseda University Grant for Special Research Projects and for The 21st Century COE Program (Holistic Research and Education Center for Physics Self-organization Systems) at Waseda University.

APPENDIX A: EUCLIDEAN ACTION

Here we present the Euclidean action. First, we calculate a scalar curvature and a Gauss-Bonnet term using the metric (2.11) as

$$R = 12\left(\frac{k-b'^2}{b^2}\right) - 8\frac{b''}{b}, \tag{A1}$$

$$\mathcal{L}_{GB} = 24\left(\frac{k-b'^2}{b^2}\right)^2 - 96\left(\frac{k-b'^2}{b^2}\right)\frac{b''}{b}. \tag{A2}$$

By using Einstein equations (2.12), we rewrite the bulk action (2.13) as

$$\begin{aligned} S_E^{\rm bulk} &= -\frac{4}{\kappa_5^2}\int_{\mathcal{M}} dx^5 \sqrt{g}\left[\Lambda + 2\frac{b''}{b} - 3\frac{k-b'^2}{b^2}\right] \\ &= -\frac{8V_4^{(k)}}{\kappa_5^2}\int_{r_1}^{r_2} dr\left[\Lambda + 2\frac{b''}{b} - 3\frac{k-b'^2}{b^2}\right]. \end{aligned} \tag{A3}$$

In a single-brane model, we just replace the integration range $[r_1, r_2]$ with $[0, r_0]$. In the second line of Eq. (A3), we put the factor 2 because an instanton is constructed from two copies of a patched manifold with region $[r_1, r_2]$ (or $[0, r_0]$).

We then calculate the surface term, i.e.

$$\begin{aligned} &K + 2\alpha\left(J - 2G^{\rho\sigma}K_{\rho\sigma}\right) \\ &= 4\left(\frac{b'}{b}\right)\left[1 - 4\alpha\left(\frac{b'^2}{b^2} - \frac{3k}{b^2}\right)\right], \end{aligned} \tag{A4}$$

where we have used $J_{\mu\nu} = -2\left(b'/b\right)^3 h_{\mu\nu}$ and $J = -8\left(b'/b\right)^3$. Using above results, we rewrite the boundary action

$$S_E^{\rm brane}=\frac{1}{\kappa_5^2}\int_{\partial\mathcal{M}_i}d^4x\sqrt{h}$$
$$\times\left\{4\left[\left(\frac{b'}{b}\right)\left\{1-4\alpha\left(\frac{b'^2}{b^2}-\frac{3k}{b^2}\right)\right\}\right]^{\pm}+\bar{\lambda}\right\}\text{(A5)}$$

Using the junction condition (2.16), (A5) gives

$$S_E^{\rm brane} = \frac{1}{\kappa_5^2}\int_{\partial\mathcal{M}_i}d^4x\sqrt{h}$$
$$\times \left[\left(\frac{b'}{b}\right)\left\{1-12\alpha\left(\frac{b'^2}{b^2}-\frac{3k}{b^2}\right)\right\}\right]^{\pm}. \text{(A6)}$$

Imposing Z_2 symmetry ($b'_-=-b'_+$), the Euclidean action for a two-brane model is rewritten as

$$S_E^{\rm brane} = \frac{2V_4^{(k)}}{\kappa_5^2}\left\{\left(\frac{b'_+}{b}\right)\left[1-12\alpha\left(\frac{b'^2_+}{b^2}-\frac{3k}{b^2}\right)\right]\right\}_{r=r_1}$$
$$-\frac{2V_4^{(k)}}{\kappa_5^2}\left\{\left(\frac{b'_-}{b}\right)\left[1-12\alpha\left(\frac{b'^2_-}{b^2}-\frac{3k}{b^2}\right)\right]\right\}_{r=r_2}. \text{(A7)}$$

For a single-brane model,

$$S_E^{\rm brane} = -\frac{2V_4^{(k)}}{\kappa_5^2}\left\{\left(\frac{b'_-}{b}\right)\left[1-12\alpha\left(\frac{b'^2_-}{b^2}-\frac{3k}{b^2}\right)\right]\right\}_{r=r_0}. \text{(A8)}$$

Finally we obtain the total action by $S_E^{\rm bulk}+S_E^{\rm brane}$.

[1] A. Vilenkin, Phys. Lett. **117B**, 25 (1982).

[2] J. B. Hartle and S. W. Hawking, Phys. Rev. D **28**, 2960 (1983).

[3] S. Coleman and F. de Luccia, Phys. Rev. D **21**, 3305 (1980).

[4] K. Akama, Lect. Notes Phys. **176**, 267 (1982).

[5] V. A. Rubakov and M. E. Shaposhnikov Phys. Lett. **152B**, 136 (1983).

[6] J. Polchinski, Phys. Rev. Lett. **75**, 4724 (1995).

[7] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. B**429**, 263 (1998); I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. B **436**, 257 (1998); N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Rev. D**59**, 086004 (1999); N. Arkani-Hamed, S. Dimopoulos, N. Kaloper, J. March-Russell, Nucl. Phys. **B567**, 189 (2000).

[8] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690 (1999).

[9] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999).

[10] P. Binétruy, adn, C. Deffayet, and Langlois, Nucl. Phys.**565**, 269 (2000).

[11] T. Shiromizu, K. Maeda, and M. Sasaki, Phys. Rev. D **62**, 024012 (2000).

[12] J. Garriga and M. Sasaki, Phys. Rev. D **62**, 043523 (2000).

[13] S. W. Hawking, T. Hertog, and H. S. Reall, Phys. Rev. D **63**, 083504 (2001).

[14] S. W. Hawking, T. Hertog, and H. S. Reall, Phys. Rev. D **62**, 043501 (2000).

[15] A. A. Strobinsky, Phys. Lett. **91B**, 99 (1980).

[16] G. Dvali, G. Gabadadze, and M. Porrati, Phys. lett. B **485**, 208 (2000); G. Dvali and G. Gabadadze, Phys. Rev. D **63**, 065007 (2001); G. Dvali, G. Gabadadze, and M. Shifman, *ibid.* **67**044020 (2003).

[17] K. Maeda, S. Mizuno, and T. Torii, Phys. Rev. D **68**, 0234033 (2003).

[18] B. Zwiebach, Phys. Lett. B **156**, 315 (1985).

[19] B. Zumino, Phys. Rep. **137**, 109 (1986).

[20] N. Deruelle and M. Sasaki, Prog.Theor.Phys. **110**, 441 (2003).

[21] J. E. Kim and H. M. Lee, Nucl. Phys. **602**, 346 (2001); *ibid.* **619**, 763 (2001).

[22] K. Maeda and T. Torii, Phys. Rev. D **69**, 024002 (2003).

[23] C. Charmousis and J. Dufaux, hep-th/0311267.

[24] M. Minamitsuji and M. Sasaki, hep-th/0404166.

[25] R. C. Myers, Phys. Rev. D **36**, 392 (1987).

[26] G. W. Gibbons and S. W. Hawking, Phys. Rev. D **15**, 2752 (1977).

[27] H. A. Chamblin and H. S. Reall, Nucl. Phys. **B562**, 133 (1999).

[28] W. Israel, Nouvo Cimento Soc. Ital. Fis. B **44**, 1 (1966).

[29] E. Witten, Nucl. Phys. **B195**, 481 (1982).

[30] N. D. Birrell and P. C. W. Davies, *Quantum fields in curved space* (Cambridge University Press, Cambridge, England, 1982).

[31] A. Vilenkin, Phys. Rev. D **32**, 2511 (1985).